

APPENDIX O
Effective Habitat Analyses

List of FiguresMiddle Fork American River – MF26.2

- Figure O-1. Middle Fork American River MF26.2 Upper Effective FYLF Egg Mass Habitat Matrix.
- Figure O-2. Middle Fork American River MF26.2 Upper Effective FYLF Tadpole Habitat Matrix.
- Figure O-3. Middle Fork American River MF26.2 Lower Effective FYLF Egg Mass Habitat Matrix.
- Figure O-4. Middle Fork American River MF26.2 Lower Effective FYLF Tadpole Habitat Matrix.

Rubicon River – R3.5

- Figure O-5. Rubicon River R3.5 Upper Effective FYLF Egg Mass Habitat Matrix.
- Figure O-6. Rubicon River R3.5 Upper Effective FYLF Tadpole Habitat Matrix.
- Figure O-7. Rubicon River R3.5 Lower Effective FYLF Egg Mass Habitat Matrix.
- Figure O-8. Rubicon River R3.5 Lower Effective FYLF Tadpole Habitat Matrix.

Middle Fork American River at Fords Bar – MF14.1*Effective Habitat Matrices*

- Figure O-9. Middle Fork American River MF14.1 Effective Rainbow Trout Spawning Habitat Matrix.
- Figure O-10. Middle Fork American River MF14.1 Effective Rainbow Trout Incubation Habitat Matrix.
- Figure O-11. Middle Fork American River MF14.1 Effective Rainbow Trout Spawning Depth Sensitivity Habitat Matrix.
- Figure O-12. Middle Fork American River MF14.1 Effective Rainbow Trout Depth Sensitivity Incubation Habitat Matrix.
- Figure O-13. Middle Fork American River MF14.1 Effective FYLF Egg Mass Habitat Matrix.
- Figure O-14. Middle Fork American River MF14.1 Effective FYLF Tadpole Habitat Matrix.
- Figure O-15. Middle Fork American River MF14.1 Effective Food Production Habitat Matrix.

Effective Habitat Time Series

- Figure O-16. Fords Bar MF14.1 Effective Habitat Time Series. (Top) Effective Rainbow Trout Spawning (Average of Eight Weekly Spawning Periods May–April) and Subsequent Incubation and (Bottom) Effective Foothill Yellow-Legged Frog Breeding Habitat (Average of Three 10–11 Day Breeding Periods (May 10–June 15) (see text for details).
- Figure O-17. Fords Bar MF14.1 Effective Rainbow Trout Spawning Habitat. Maximum, Minimum, and 25% and 75% Exceedance of Eight Weekly Spawning Periods (March–April) and Subsequent Incubation (see text for details).

Middle Fork American River at Buckeye Bar – MF4.8

Effective Habitat Matrices

- Figure O-18. Middle Fork American River MF4.8 Effective Rainbow Trout Spawning Habitat Matrix.
- Figure O-19. Middle Fork American River MF4.8 Effective Rainbow Trout Incubation Habitat Matrix.
- Figure O-20. Middle Fork American River MF4.8 Effective Rainbow Trout Spawning Depth Sensitivity Habitat Matrix.
- Figure O-21. Middle Fork American River MF4.8 Effective Rainbow Trout Depth Sensitivity Incubation Habitat Matrix.
- Figure O-22. Middle Fork American River MF4.8 Effective FYLF Egg Mass Habitat Matrix.
- Figure O-23. Middle Fork American River MF4.8 Effective FYLF Tadpole Habitat Matrix.
- Figure O-24. Middle Fork American River MF4.8 Effective Food Production Habitat Matrix.

Effective Habitat Time Series

- Figure O-25. Buckeye Bar MF4.8 Effective Habitat Time Series. (Top) Effective Rainbow Trout Spawning Habitat (Average of Eight Weekly Spawning Periods March–April) and Subsequent Incubation and (Bottom) Effective Foothill Yellow-Legged Frog Breeding Habitat (Average of Three 10–11 Day Breeding Periods (May 10–June 15) (see text for details).
- Figure O-26. Buckeye Bar MF4.8 Effective Rainbow Trout Spawning Habitat. Maximum, Minimum, and 25% and 75% Exceedance of Eight Weekly Spawning Periods (March–April) and Subsequent Incubation (see text for details).

FIGURES